

Search Title: 00-1041b.opt User: cpamoj - < Not Set >, < Not Set >  
 PAN: 00-056721, Page 1 of 2, Tue Oct 10 10:20:11, VIEWED MARKED

<p>2000-056721/05 B02 D13 HAYB 1998.04.20          HAYASHIBARA SEIBUTSU KAGAKU          1998.04.20 1998-109393(+1998JP-109393) (1999.11.02) A23L 1/30,          1/06, 2/52 // A61K 31/35, C07D 311/62          Nourishment liquid useful for preparing jelly, soft drink,          foodstuffs, etc and also for preventing geriatric diseases,          antioxdation in blood - consists of proanthocyanidine as          antioxidant          C2000-015405          Addtl. Data: HOMETTO KK (HOME-)</p>	<p>B(6-A1, 14-F2B, 14-F2D, 14-F4, 14-F6, 14-N3, 14-          N17C, 14-S4, 14-S8) D(3-H1J, 3-H1T2) .4          agar jelly (claimed). Clinical applications of the foodstuff or drink          includes rehabilitation of diabetes mellitus, hypertension,          hyperlipidemia, atopic dermatitis and cerebral infarction. Also          improves micro-circulation of blood, platelet aggregation, vision and          prevents ageing.</p>
<p><u>NOVELTY</u>          The nourishment liquid is an antioxdation liquid formulation          containing proanthocyanidine as active ingredient which is stable in          aqueous solution.</p>	<p><u>ADVANTAGE</u>          The nourishment liquid has sweet taste, heat resistance, anticaries,          non-coloring property and reduced astringency. It has a high          nutritional value and is easily digested and absorbed by the small          intestine.</p>
<p><u>DETAILED DESCRIPTION</u>          An INDEPENDENT CLAIM is also included for manufacturing          method of proanthocyanidine containing nourishment liquid,          comprising dissolving proanthocyanidine and trehalose in water.</p>	<p><u>ORGANIC CHEMISTRY</u>          Preferred composition: To the nourishment liquid, trehalose is          added as an adjuvant for providing stabilizing effect.</p>
<p><u>USE</u>          The nourishment liquid is used for preparing foodstuffs, drink and          and a drink was prepared. Color, transparency, pH and SOD activity          was tested. The SOD activity in 40 °C was found to be reduced to 60-          70% in a month. Change in pH, turbidity and color was not observed.          (RP2)          (14pp3119DWgNo.0/7)</p>	<p><u>EXAMPLE</u>          Trehalose and polyphenol were mixed with proanthocyanidine</p>

JP 11299452-A